

Mathieu groups as Galois groups

Peter Müller

LMU, 10 June 2015

The inverse Galois problem

G finite group, is there $f(X) \in \mathbb{Q}[X]$ with $\text{Gal}(f(X)/\mathbb{Q}) = G$?

The inverse Galois problem

G finite group, is there $f(X) \in \mathbb{Q}[X]$ with $\text{Gal}(f(X)/\mathbb{Q}) = G$?

Yes, if

- G abelian (algebra class)
- G solvable (Shafarevich + ε)
- G one of the 26 simple sporadic groups, except possibly M_{23}
- many more cases (mainly by rigidity)
- G arbitrary?

The inverse Galois problem

G finite group, is there $f(X) \in \mathbb{Q}[X]$ with $\text{Gal}(f(X)/\mathbb{Q}) = G$?

Yes, if

- G abelian (algebra class)
- G solvable (Shafarevich + ε)
- G one of the 26 simple sporadic groups, except possibly M_{23}
- many more cases (mainly by rigidity)
- G arbitrary?

Different base fields

Every finite group is a Galois group over $\mathbb{C}(t)$.

M_{23} as Galois group over $\mathbb{F}_2(t)$

Theorem (Abhyankar 1993)

$$f(X) = X^{23} + tX^3 + 1 \in \mathbb{F}_2(t)[X]$$

$$G = \text{Gal}(f(X)/\mathbb{F}_2(t)) = M_{23}$$

M_{23} as Galois group over $\mathbb{F}_2(t)$

Theorem (Abhyankar 1993)

$$f(X) = X^{23} + tX^3 + 1 \in \mathbb{F}_2(t)[X]$$

$$G = \text{Gal}(f(X)/\mathbb{F}_2(t)) = M_{23}$$

Proof.

- $f(X)$ irreducible $\implies G$ transitive.

M_{23} as Galois group over $\mathbb{F}_2(t)$

Theorem (Abhyankar 1993)

$$f(X) = X^{23} + tX^3 + 1 \in \mathbb{F}_2(t)[X]$$
$$G = \text{Gal}(f(X)/\mathbb{F}_2(t)) = M_{23}$$

Proof.

- $f(X)$ irreducible $\implies G$ transitive.
- $X^{23} + X^3 + 1 = (X^3 + X^2 + 1)(X^5 + \dots)(X^{15} + \dots) \implies G$ has elements of order 15 $\implies G = M_{23}, A_{23}$ or S_{23} .

M_{23} as Galois group over $\mathbb{F}_2(t)$

Theorem (Abhyankar 1993)

$$f(X) = X^{23} + tX^3 + 1 \in \mathbb{F}_2(t)[X]$$
$$G = \text{Gal}(f(X)/\mathbb{F}_2(t)) = M_{23}$$

Proof.

- $f(X)$ irreducible $\implies G$ transitive.
- $X^{23} + X^3 + 1 = (X^3 + X^2 + 1)(X^5 + \dots)(X^{15} + \dots) \implies G$ has elements of order 15 $\implies G = M_{23}, A_{23}$ or S_{23} .
- Serre's linearization trick: $f(X)$ divides **additive polynomial**

$$\hat{f}(X) = X^{2^{11}} + t^{16}X^{2^8} + \dots + tX^{2^2} + t^8X^2 + X, \text{ hence}$$

$$\text{GL}_{11}(\mathbb{F}_2) \geq \text{Gal}(\hat{f}(X)/\mathbb{F}_2(t)) \twoheadrightarrow G \implies 19 \nmid |G| \implies A_{23} \not\leq G.$$



Serre's linearization trick

Serre's trick works ...

... because \mathbb{F}_2 -permutation module of M_{23} has 11-dimensional submodule (even words of Golay code).

Similar trick works over \mathbb{F}_3 for $M_{11} \leq S_{12}$.

Serre's linearization trick

Serre's trick works ...

... because \mathbb{F}_2 -permutation module of M_{23} has 11-dimensional submodule (even words of Golay code).

Similar trick works over \mathbb{F}_3 for $M_{11} \leq S_{12}$.

Conway, McKay, Trojan (2009)

$$X^{23} + X^3 + t \quad \mathbb{F}_2(t) \quad M_{23}$$

$$X^{24} + X + t \quad \mathbb{F}_2(t) \quad M_{24}$$

Serre's linearization trick

Serre's trick works ...

... because \mathbb{F}_2 -permutation module of M_{23} has 11-dimensional submodule (even words of Golay code).

Similar trick works over \mathbb{F}_3 for $M_{11} \leq S_{12}$.

Conway, McKay, Trojan (2009)

$X^{23} + X^3 + t$	$\mathbb{F}_2(t)$	M_{23}
$X^{24} + X + t$	$\mathbb{F}_2(t)$	M_{24}
$X^{12} + X + t$	$\mathbb{F}_3(t)$	M_{11}

Serre's linearization trick

Serre's trick works ...

... because \mathbb{F}_2 -permutation module of M_{23} has 11-dimensional submodule (even words of Golay code).

Similar trick works over \mathbb{F}_3 for $M_{11} \leq S_{12}$.

Conway, McKay, Trojan (2009)

$X^{23} + X^3 + t$	$\mathbb{F}_2(t)$	M_{23}
$X^{24} + X + t$	$\mathbb{F}_2(t)$	M_{24}
$X^{12} + X + t$	$\mathbb{F}_3(t)$	M_{11}
$X^{11} + tX^2 - 1$	$\mathbb{F}_3(t)$	M_{11}

Serre's linearization trick

Serre's trick works ...

... because \mathbb{F}_2 -permutation module of M_{23} has 11-dimensional submodule (even words of Golay code).

Similar trick works over \mathbb{F}_3 for $M_{11} \leq S_{12}$.

Conway, McKay, Trojan (2009)

$X^{23} + X^3 + t$	$\mathbb{F}_2(t)$	M_{23}
$X^{24} + X + t$	$\mathbb{F}_2(t)$	M_{24}
$X^{12} + X + t$	$\mathbb{F}_3(t)$	M_{11}
$X^{11} + tX^2 - 1$	$\mathbb{F}_3(t)$	M_{11}

Bad characteristic

How to treat M_{23} in characteristic $\neq 2$?

Mathieu group M_{23} as monodromy group of a polynomial

Properties of M_{23}

- $M_{23} \leq S_{23}$ is 4-transitive on $\{1, 2, \dots, 23\}$.
- $|M_{23}| = 23 \cdot 22 \cdot 21 \cdot 20 \cdot 48 = 10200960$.
- $[A_{23} : M_{23}] = 1267136462592000$.
- M_{23} is simple.
- M_{23} is self-normalizing in S_{23} .

Mathieu group M_{23} as monodromy group of a polynomial

Properties of M_{23}

- $M_{23} \leq S_{23}$ is 4-transitive on $\{1, 2, \dots, 23\}$.
- $|M_{23}| = 23 \cdot 22 \cdot 21 \cdot 20 \cdot 48 = 10200960$.
- $[A_{23} : M_{23}] = 1267136462592000$.
- M_{23} is simple.
- M_{23} is self-normalizing in S_{23} .

M_{23} as monodromy group of a polynomial

- $\hat{h}(X) \in \mathbb{C}[X]$, such that $\text{Gal}(\hat{h}(X) - t/\mathbb{C}(t)) = M_{23}$,
- (equivalent to) $\text{Mon}(\mathbb{P}^1(\mathbb{C}) \rightarrow \mathbb{P}^1(\mathbb{C}), z \mapsto \hat{h}(z)) = M_{23}$.
- Existence: Well known and easy (up to Riemann's existence theorem),
- with unique branching type: $1^7 2^8, 1^3 2^2 4^4, 23^1$.

Mathieu group M_{23} as monodromy group of a polynomial

Theorem (Matiyasevich 1998 (unpublished), Elkies 2013)

$$K = \mathbb{Q}(\sqrt{23 - 2\sqrt{-23}})$$

$\hat{h}(X) = X^{23} + \text{complicated lower order terms} \in K[X]:$

$$\text{Gal}(\hat{h}(X) - t/K(t)) = M_{23}.$$

Mathieu group M_{23} as monodromy group of a polynomial

Theorem (Matiyasevich 1998 (unpublished), Elkies 2013)

$$K = \mathbb{Q}(\sqrt{23 - 2\sqrt{-23}})$$

$\hat{h}(X) = X^{23} + \text{complicated lower order terms} \in K[X]$:

$$\text{Gal}(\hat{h}(X) - t/K(t)) = M_{23}.$$

Analytic verification of Galois group

Numerically compute monodromy group of cover $\mathbb{P}^1(\mathbb{C}) \rightarrow \mathbb{P}^1(\mathbb{C})$
 $z \mapsto \hat{h}(z)$.

Mathieu group M_{23} as monodromy group of a polynomial

Theorem (Matiyasevich 1998 (unpublished), Elkies 2013)

$$K = \mathbb{Q}(\sqrt{23 - 2\sqrt{-23}})$$

$\hat{h}(X) = X^{23} + \text{complicated lower order terms} \in K[X]$:

$$\text{Gal}(\hat{h}(X) - t/K(t)) = M_{23}.$$

Analytic verification of Galois group

Numerically compute monodromy group of cover $\mathbb{P}^1(\mathbb{C}) \rightarrow \mathbb{P}^1(\mathbb{C})$
 $z \mapsto \hat{h}(z)$.

Algebraic verification of Galois group, first step

- Pick prime $p > 23$, such that $h(X) = (\hat{h}(X) \bmod p) \in \mathbb{F}_p[X]$.
- Suffices to show: $\text{Gal}(h(X) - t/\mathbb{F}_p(t)) = M_{23}$. (S. Beckmann)
- Easy: $M_{23} \leq \text{Gal}(h(X) - t/\mathbb{F}_p(t))$. (Dedekind)
- Need to decide: $\text{Gal}(h(X) - t/\mathbb{F}_p(t)) = M_{23}$ or A_{23} ?

Verification of Galois group

Naive idea, for small $p > 23$:

- M_{23} has two orbits on 5-sets, of lengths 5313 and 28336.

Verification of Galois group

Naive idea, for small $p > 23$:

- M_{23} has two orbits on 5-sets, of lengths 5313 and 28336.
- “Compute” polynomial of degree $\binom{23}{5} = 33649$, whose roots are the 5-sums of roots of $h(X) - t$, and “check” if it has a degree 5313 factor over $\mathbb{F}_p(t) \dots$

Verification of Galois group

Naive idea, for small $p > 23$:

- M_{23} has two orbits on 5-sets, of lengths 5313 and 28336.
- “Compute” polynomial of degree $\binom{23}{5} = 33649$, whose roots are the 5-sums of roots of $h(X) - t$, and “check” if it has a degree 5313 factor over $\mathbb{F}_p(t) \dots$

Using Weil-bound for points on curves (Elkies)

$$\frac{1}{|G|} = \lim_{p \rightarrow \infty} \frac{|\{t_0 \in \mathbb{F}_p \mid h(X) - t_0 \text{ splits into linear factors}\}|}{p}$$

Elkies chose $p = 10^8 + 7$: *The factorization of 10^8 polynomials mod p was a somewhat extravagant computation (two days of CPU time in gp).*

M_{23} and its Steiner system

Steiner system $S = S(4, 7, 23)$

- $\mathfrak{P} = 23$ points, $\mathfrak{B} = 253$ blocks = certain 7-sets from \mathfrak{P}
- $|\mathfrak{B}| \binom{7}{4} = \binom{23}{4}$ (any 4-set of points contained in exactly one block)

M_{23} and its Steiner system

Steiner system $S = S(4, 7, 23)$

- $\mathfrak{P} = 23$ points, $\mathfrak{B} = 253$ blocks = certain 7-sets from \mathfrak{P}
- $|\mathfrak{B}| \binom{7}{4} = \binom{23}{4}$ (any 4-set of points contained in exactly one block)
- $M_{23} = \text{Aut}(S)$ transitive on \mathfrak{P} and \mathfrak{B}

M_{23} and its Steiner system

Steiner system $S = S(4, 7, 23)$

- $\mathfrak{P} = 23$ points, $\mathfrak{B} = 253$ blocks = certain 7-sets from \mathfrak{P}
- $|\mathfrak{B}| \binom{7}{4} = \binom{23}{4}$ (any 4-set of points contained in exactly one block)
- $M_{23} = \text{Aut}(S)$ transitive on \mathfrak{P} and \mathfrak{B}
- $h(X) - t = \prod_{x \in \mathfrak{P}} (X - x)$. Fix $x_0 \in \mathfrak{P}$, so $t = h(x_0)$.

M_{23} and its Steiner system

Steiner system $S = S(4, 7, 23)$

- $\mathfrak{P} = 23$ points, $\mathfrak{B} = 253$ blocks = certain 7-sets from \mathfrak{P}
- $|\mathfrak{B}| \binom{7}{4} = \binom{23}{4}$ (any 4-set of points contained in exactly one block)
- $M_{23} = \text{Aut}(S)$ transitive on \mathfrak{P} and \mathfrak{B}
- $h(X) - t = \prod_{x \in \mathfrak{P}} (X - x)$. Fix $x_0 \in \mathfrak{P}$, so $t = h(x_0)$.

Associated polynomials

$x \in \mathfrak{P}$ integral over $\mathbb{F}_p[t]$ and $\mathbb{F}_p[x_0]$, hence

$$H(h(x_0), Y) = H(t, Y) = \prod_{B \in \mathfrak{B}} (Y - \sum_{x \in B} x) \in \mathbb{F}_p[t][Y] \quad (\text{degree } 253)$$

M_{23} and its Steiner system

Steiner system $S = S(4, 7, 23)$

- $\mathfrak{P} = 23$ points, $\mathfrak{B} = 253$ blocks = certain 7-sets from \mathfrak{P}
- $|\mathfrak{B}| \binom{7}{4} = \binom{23}{4}$ (any 4-set of points contained in exactly one block)
- $M_{23} = \text{Aut}(S)$ transitive on \mathfrak{P} and \mathfrak{B}
- $h(X) - t = \prod_{x \in \mathfrak{P}} (X - x)$. Fix $x_0 \in \mathfrak{P}$, so $t = h(x_0)$.

Associated polynomials

$x \in \mathfrak{P}$ integral over $\mathbb{F}_p[t]$ and $\mathbb{F}_p[x_0]$, hence

$$H(h(x_0), Y) = H(t, Y) = \prod_{B \in \mathfrak{B}} (Y - \sum_{x \in B} x) \in \mathbb{F}_p[t][Y] \quad (\text{degree } 253)$$

$$H_1(x_0, Y) = \prod_{x_0 \in B \in \mathfrak{B}} (Y - \sum_{x \in B} x) \in \mathbb{F}_p[x_0][Y] \quad (\text{degree } 77)$$

M_{23} and its Steiner system

Steiner system $S = S(4, 7, 23)$

- $\mathfrak{P} = 23$ points, $\mathfrak{B} = 253$ blocks = certain 7-sets from \mathfrak{P}
- $|\mathfrak{B}| \binom{7}{4} = \binom{23}{4}$ (any 4-set of points contained in exactly one block)
- $M_{23} = \text{Aut}(S)$ transitive on \mathfrak{P} and \mathfrak{B}
- $h(X) - t = \prod_{x \in \mathfrak{P}} (X - x)$. Fix $x_0 \in \mathfrak{P}$, so $t = h(x_0)$.

Associated polynomials

$x \in \mathfrak{P}$ integral over $\mathbb{F}_p[t]$ and $\mathbb{F}_p[x_0]$, hence

$$H(h(x_0), Y) = H(t, Y) = \prod_{B \in \mathfrak{B}} (Y - \sum_{x \in B} x) \in \mathbb{F}_p[t][Y] \quad (\text{degree } 253)$$

$$H_1(x_0, Y) = \prod_{x_0 \in B \in \mathfrak{B}} (Y - \sum_{x \in B} x) \in \mathbb{F}_p[x_0][Y] \quad (\text{degree } 77)$$

$$H_2(x_0, Y) = \prod_{x_0 \notin B \in \mathfrak{B}} (Y - \sum_{x \in B} x) \in \mathbb{F}_p[x_0][Y] \quad (\text{degree } 176)$$

Lemma (essentially)

If $h(X) \in \mathbb{F}_p[X]$ has degree 23, then

$$\text{Gal}(h(X) - t/\mathbb{F}_p(t)) = M_{23} \iff H(h(X), Y) = H_1(X, Y)H_2(X, Y)$$

for some $H(t, Y) \in \mathbb{F}_p[t][Y]$ irreducible of degree 253, and $H_1(X, Y), H_2(X, Y) \in \mathbb{F}_p[X][Y]$ of degrees 77 and 176.

Lemma (essentially)

If $h(X) \in \mathbb{F}_p[X]$ has degree 23, then

$$\text{Gal}(h(X) - t/\mathbb{F}_p(t)) = M_{23} \iff H(h(X), Y) = H_1(X, Y)H_2(X, Y)$$

for some $H(t, Y) \in \mathbb{F}_p[t][Y]$ irreducible of degree 253, and $H_1(X, Y), H_2(X, Y) \in \mathbb{F}_p[X][Y]$ of degrees 77 and 176.

How to compute ...

$$H(t, Y) = \prod_{B \in \mathfrak{B}} (Y - \sum_{x \in B} x)$$

from $h(X)$? Certainly not as a degree 253 factor of the degree $\binom{23}{7} = 245157$ polynomial

$$\prod_{C \in \binom{\mathfrak{B}}{7}} (Y - \sum_{x \in C} x).$$

Laurent series

Computation of ...

$$H(t, Y) = \prod_{B \in \mathfrak{B}} (Y - \sum_{x \in B} x) \in \mathbb{F}_p[t][Y] = \mathbb{F}_p[\tau^{23}][Y]$$

by **explicit** determination of \mathfrak{P} and \mathfrak{B} :

Laurent series

Computation of ...

$$H(t, Y) = \prod_{B \in \mathfrak{B}} (Y - \sum_{x \in B} x) \in \mathbb{F}_p[t][Y] = \mathbb{F}_p[\tau^{23}][Y]$$

by **explicit** determination of \mathfrak{P} and \mathfrak{B} :

- $h(X) - t = h(X) - \tau^{23} = 0$ has a root

$$L(\tau) = \tau + a_0 + a_1\tau^{-1} + a_2\tau^{-2} + \cdots \in \mathbb{F}_p((1/\tau)).$$

Laurent series

Computation of ...

$$H(t, Y) = \prod_{B \in \mathfrak{B}} (Y - \sum_{x \in B} x) \in \mathbb{F}_p[t][Y] = \mathbb{F}_p[\tau^{23}][Y]$$

by **explicit** determination of \mathfrak{P} and \mathfrak{B} :

- $h(X) - t = h(X) - \tau^{23} = 0$ has a root

$$L(\tau) = \tau + a_0 + a_1\tau^{-1} + a_2\tau^{-2} + \cdots \in \mathbb{F}_p((1/\tau)).$$

- $\mathfrak{P} = \{L(w\tau) \mid w \in W\}$ where $W \leq \bar{\mathbb{F}}_p^*$ with $|W| = 23$.

Laurent series

Computation of ...

$$H(t, Y) = \prod_{B \in \mathfrak{B}} (Y - \sum_{x \in B} x) \in \mathbb{F}_p[t][Y] = \mathbb{F}_p[\tau^{23}][Y]$$

by **explicit** determination of \mathfrak{P} and \mathfrak{B} :

- $h(X) - t = h(X) - \tau^{23} = 0$ has a root

$$L(\tau) = \tau + a_0 + a_1\tau^{-1} + a_2\tau^{-2} + \cdots \in \mathbb{F}_p((1/\tau)).$$

- $\mathfrak{P} = \{L(w\tau) \mid w \in W\}$ where $W \leq \bar{\mathbb{F}}_p^*$ with $|W| = 23$.
- W acts regularly on \mathfrak{P} .

Laurent series

Computation of ...

$$H(t, Y) = \prod_{B \in \mathfrak{B}} (Y - \sum_{x \in B} x) \in \mathbb{F}_p[t][Y] = \mathbb{F}_p[\tau^{23}][Y]$$

by **explicit** determination of \mathfrak{P} and \mathfrak{B} :

- $h(X) - t = h(X) - \tau^{23} = 0$ has a root

$$L(\tau) = \tau + a_0 + a_1\tau^{-1} + a_2\tau^{-2} + \dots \in \mathbb{F}_p((1/\tau)).$$

- $\mathfrak{P} = \{L(w\tau) \mid w \in W\}$ where $W \leq \bar{\mathbb{F}}_p^*$ with $|W| = 23$.
- W acts regularly on \mathfrak{P} .
- There are only **two** continuations of W to an action of M_{23} , so there are only **two** candidates for \mathfrak{B} . One of them works!

Laurent series

Computation of ...

$$H(t, Y) = \prod_{B \in \mathfrak{B}} (Y - \sum_{x \in B} x) \in \mathbb{F}_p[t][Y] = \mathbb{F}_p[\tau^{23}][Y]$$

by **explicit** determination of \mathfrak{P} and \mathfrak{B} :

- $h(X) - t = h(X) - \tau^{23} = 0$ has a root

$$L(\tau) = \tau + a_0 + a_1\tau^{-1} + a_2\tau^{-2} + \dots \in \mathbb{F}_p((1/\tau)).$$

- $\mathfrak{P} = \{L(w\tau) \mid w \in W\}$ where $W \leq \bar{\mathbb{F}}_p^*$ with $|W| = 23$.
- W acts regularly on \mathfrak{P} .
- There are only **two** continuations of W to an action of M_{23} , so there are only **two** candidates for \mathfrak{B} . One of them works!
- Suffices to work with **truncated** Laurent series.

Laurent series

Computation of ...

$$H(t, Y) = \prod_{B \in \mathfrak{B}} (Y - \sum_{x \in B} x) \in \mathbb{F}_p[t][Y] = \mathbb{F}_p[\tau^{23}][Y]$$

by **explicit** determination of \mathfrak{P} and \mathfrak{B} :

- $h(X) - t = h(X) - \tau^{23} = 0$ has a root

$$L(\tau) = \tau + a_0 + a_1\tau^{-1} + a_2\tau^{-2} + \dots \in \mathbb{F}_p((1/\tau)).$$

- $\mathfrak{P} = \{L(w\tau) \mid w \in W\}$ where $W \leq \bar{\mathbb{F}}_p^*$ with $|W| = 23$.
- W acts regularly on \mathfrak{P} .
- There are only **two** continuations of W to an action of M_{23} , so there are only **two** candidates for \mathfrak{B} . One of them works!
- Suffices to work with **truncated** Laurent series.
- **No need** to factor $H(h(X), Y)$ to obtain $H_1(X, Y)$ and $H_2(X, Y)$.
Work in $\bar{\mathbb{F}}_p((1/x_0))!$

Laurent series

More general case

Want to upper bound $G = \text{Gal}(P(X) - tQ(X)/k(t)) \leq S_n$. Method works best,

- if there is a set B with $2 \leq |B| \leq n - 2$, and $[G : G_B]$ small,
- there is an inertia generator with few cycles (hence few potential candidates for B), and
- k is a finite field (otherwise the coefficients of Laurent series explode).

For instance, it works well for Granboulan's M_{24} -polynomial.

Reverting the technique to find polynomials

Using $H_1(x_0, Y)$

$$\text{Gal}(h(X) - t/\mathbb{F}_p(t)) = M_{23} \implies \prod_{x_0 \in B \in \mathfrak{B}} (Y - \sum_{x \in B} x) \in \mathbb{F}_p[x_0][Y],$$

hence

$$S_k = \sum_{x_0 \in B \in \mathfrak{B}} (\sum_{x \in B} x)^k \in \mathbb{F}_p[x_0] \text{ for all } k \geq 0.$$

Reverting the technique to find polynomials

Using $H_1(x_0, Y)$

$$\text{Gal}(h(X) - t/\mathbb{F}_p(t)) = M_{23} \implies \prod_{x_0 \in B \in \mathfrak{B}} (Y - \sum_{x \in B} x) \in \mathbb{F}_p[x_0][Y],$$

hence

$$S_k = \sum_{x_0 \in B \in \mathfrak{B}} (\sum_{x \in B} x)^k \in \mathbb{F}_p[x_0] \text{ for all } k \geq 0.$$

On the other hand, with $x_0 = 1/z$, ω a 23-rd root of unity, and $m > 0$, the roots x_i of $h(X) - t = h(X) - h(1/z)$ are

$$x_i = \frac{\omega^i}{z} + \text{higher order terms} = A_i(z) + O(z^m) \in \mathbb{F}_p((z)), \text{ hence}$$

$$\mathbb{F}_p[1/z] \ni S_k = \sum_{x_0 \in B \in \mathfrak{B}} (\sum_{x_i \in B} A_i(z))^k + O(z^{m+1-k}).$$

Reverting the technique to find polynomials

Strategy:

- Set $h(X) = a_1X + a_2X^2 + \cdots + a_{21}X^{21} + X^{23} \in \mathbb{F}_p[\mathbf{a}][X]$.
- For $m > 0$ compute $x_i = A_i(z) + O(z^m) \in \mathbb{F}_p[\mathbf{a}]((z))$.
- For $k = 1, 2, \dots, m-1$ collect the coefficients of z^j with $j \geq 1$ in $\sum_{x_0 \in B \in \mathfrak{B}} (\sum_{x \in B} A_i(z))^k$. They all have to vanish!
- Solve this system of polynomial equations for the unknowns \mathbf{a} .

Results:

- For $p = 47$ get Elkies' polynomial within a few seconds (compared to 46 CPU hours by refined standard approach).
- One can also compute the Laurent series and Gröbner bases over \mathbb{Q} instead of \mathbb{F}_p . Then a naive Sage implementation takes a few minutes to get the degree 4 number field over which the polynomial is defined.